

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/40 // (A61K 31/40, 31:00, 38:00)		A1	(11) International Publication Number: WO 95/28926 (43) International Publication Date: 2 November 1995 (02.11.95)
(21) International Application Number: PCT/IB95/00332 (22) International Filing Date: 21 April 1995 (21.04.95) (30) Priority Data: 08/231,969 21 April 1994 (21.04.94) US 08/253,938 2 June 1994 (02.06.94) US 08/327,478 21 October 1994 (21.10.94) US 08/340,416 15 November 1994 (15.11.94) US (71) Applicant: SCHERING AKTIENGESELLSCHAFT [DE/DE]; Berlin-Wedding, Muellerstrasse 178, D-13342 Berlin (DE). (71)(72) Applicant and Inventor: PEREZ, H., Daniel [US/US]; 215 McAllister Avenue, Kentfield, CA 94904 (US). (72) Inventors: WACHTEL, Helmut; Suarezstrasse 22, D-14057 Berlin (DE). LÖSCHMANN, Peter-Andreas; Leon- Schmalzbach-Weg 7, D-7450 Hechingen (DE). GRAF, Hermann; Olwenstrasse 62, D-13465 Berlin (DE). HEDG- PETH, Joel; 33 Robinhood, San Francisco, CA 94127 (US). (74) Agent: HEANEY, Brion, P.; Millen, White, Zelano & Brani- gan, P.C., Arlington Courthouse Plaza 1, Suite 1400, 2200 Clarendon Boulevard, Arlington, VA 22201 (US).			(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PDE IV INHIBITORS FOR TREATING MULTIPLE SCLEROSIS			
(57) Abstract The present invention is a method of preventing or ameliorating the episodic recurrence of MS, comprising administering an effective amount of selective phosphodiesterase inhibitors of Type IV, e.g., Rolipram, e.g., wherein the severity of the episodic recurrences is ameliorated or the time period between episodes is lengthened. The present invention also relates to a pharmaceutical composition for treating MS comprising an effective amount of a combination of a PDE IV inhibitor and an anti-inflammatory or immunomodulatory drug in a pharmaceutically acceptable carrier.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

PDE IV inhibitors for treating multiple sclerosis

Background of the Invention

Demyelinating diseases are severe afflictions of the brain and spinal cord, involving the destruction of the myelin sheath which surrounds nerve fibers. As a result of demyelination, various neurological symptoms are manifested, including motor impairment, visual loss, and sensory changes. Multiple sclerosis (MS) is the most common of the demyelinating diseases. It is a disease characterized by episodes of focal disorder of the optic nerves, spinal cord, and brain. It is a severe chronic disabling disease with characteristic demyelination in the CNS triggered by probable autoimmune mechanisms in a genetically susceptible population. Typically, it produces recurring episodes of neurologic dysfunction, followed by remission (relapsing-remitting), but it may also be chronic. Although MS is a well-studied disease, its precise cause remains undetermined. One explanation is that an environmental factor which most often acts in childhood activates a specific population of T-cells (with the potential of attacking myelin-associated antigens such as MBP, MAP, MOP or others) which normally is controlled by suppressor-cells. In many MS cases, non-specific stress results in disease exacerbations with opening of the BBB, edema, immigration and activation of T-cells and macrophages and subsequent destruction of oligodendroglia- associated myelin followed by failing attempts at remyelination and finally a glia scar (relapsing - remitting form of MS). These exacerbations which can be visualized by MRI are associated, depending on location, with severe functional disabilities;

- 2 -

furthermore, with an increasing number of relapses, the disease (and the antigen being attacked) becomes more generalized and progresses with less clearcut intervals (progressive form of MS, secondary or primary) and increasing and persisting disability which causes (in this mostly young population) impairment of life quality with loss of employment and independent life (with hospitalization and eventually death).

Until recently, treatments have been empirical and not entirely successful. See Cecil's Textbook of Medicine (Wyngaarden, 1993). Classic non-specific immunosuppressive therapies (including corticosteroids) and cytostatic drugs as a rule have failed to alter this sequence of events and disease progression - possibly because they are poorly tolerated in these patients and also because they inhibit endogenous immunosuppressive mechanisms as well. For a review of treatments, e.g., see *Principles of Neurology*, Fifth Edition, (Adams et al., 1993). Recently non-specific immunomodulatory therapy with interferon- β -1b has been shown to prolong disease-free intervals in patients with beginning relapsing/remitting MS; however, in the great majority of these patients, exacerbations cannot be prevented completely, and also the effect on disability during the first three years of therapy is still small. Corticosteroid therapy relieves some acute symptoms of these exacerbations (probably by reducing edema) but does not affect long term prognosis. However, it appears of utmost importance to suppress all exacerbations with their activation, extension and amplification of autoimmune mechanisms which make the disease uncontrollable and disabling. There remains a need for additional drugs which have an effect on the diseases's severity and progression.

Summary of the Invention

Recent advance in the clinical, biochemical and imaging technologies enable clinicians to predict and diagnose such exacerbations in a very early stage, and thus, it now becomes

- 3 -

possible to combine different therapeutic strategies to achieve a maximal therapeutic effect. Immunomodulatory therapies with non-specific mechanisms, e.g., with interferon- β -1b (or specific immunomodulatory drugs such as copolymer I, as well as inducing tolerance to MS antigens) which will never work to 100% as they need to be given before the onset of the disease, but they can reduce to a clear and significant degree the number of clinical exacerbations and, even more, of lesions in the brain. Fortunately, for the target of bringing disease activity to a complete hold, the new diagnostic techniques being developed have the potential to detect the very early beginning of an exacerbation (e.g., increases in γ -interferons, TNF- α , increased number of activated specific T-cells in the blood, new specific MRI and other imaging techniques, and clinical observation). Therefore, strong and efficient drugs are necessary which act not just on some symptoms of the exacerbation but are able to prevent them completely and thus, inhibit disease progression. However, also new therapeutic strategies which aim, e.g., at peripheral T-cell activation, endothelial adhesion, opening of blood-brain barrier and activation of the T-cell-macrophage/-microglia interaction with subsequent oligodendrocyte damage and demyelination do affect defense mechanisms not just in the autoimmune condition but, on chronic use, are also damaging vital defense mechanism against exogenous (e.g., bacterial, parasitic or viral infections) as well as against endogenous harmful agents and effects (e.g., tumorigenesis).

An aspect of the present invention is to combine well tolerated chronic maintenance therapy with the use of new diagnostic techniques to predict or detect early exacerbations which can then, but only then, be treated aggressively by a number of drugs, or their combinations, to ensure that no persistent CNS lesions can be produced with their fatal influence on disease exacerbation, extension and progression. Furthermore, combination of therapies with different mechanisms to achieve maximum efficacy will improve tolerability of therapy (as the effects of interferon- β -1b in

- 4 -

MS have been shown to be dose-dependent, higher dosages could be expected to achieve higher and even 100% efficacy, but increasingly severe side effects prevent this type of treatment), and finally, these new combinations of maintenance and anti-exacerbation therapies result in a clearly reduced risk of side effects which can be caused by high-dose and long-term use of these drugs in monotherapy. Indeed when basic maintenance therapy is being combined with the use of these new strategies only when needed (e.g., during or just before an exacerbation) less side effects of both complementary and synergistic therapeutic lines do occur and these forms of therapy can be combined to achieve an optimal clinical result. Also, in this new concept of short-term therapy of exacerbations, different drugs can be combined to prevent further damage and disease progression. Thus, inhibition of synthesis of cytotoxic cyto- and chemokines can be combined with drugs which inhibit their release, and both may be enhanced in efficacy by, e.g., simultaneous inhibition of traffic across the blood-brain barrier, or with other drugs which inhibit - or even reverse - the ultimate tissue damage (e.g., nerve growth factors, calcipotriols, calpain inhibitors, etc.).

- 5 -

pathogenic cascade

possible therapeutic intervention

- | | | |
|-----|--|---|
| 1.- | activation of peripheral specific T-cells which are potentially autoreactive to myelin | destroy these T-cells
reactivate tolerance by oral antigens, immune globulins? enhance suppressor mechanisms, e.g., interferon- β -1b |
| 2.- | enhanced blood-brain barrier permeability to these T-cells | close blood-brain barrier, e.g., by anti-adhesion molecules such as anti-integrin monoclonal antibodies |
| 3.- | immigration of macrophages, and macrophage-T-cell-interaction and activation | interfere with T-cell receptors, enhance suppressor mechanisms, e.g., by interferon β -1b |
| 4.- | local inflammation and edema | non-specific anti-inflammatory and anti-edema strategies, e.g., corticosteroids |
| 5 | 5.- enhance release of γ -interferons, TNF α and other cytotoxic cytokines | antagonize γ -interferon, TNF α and other cytotoxic cytokines by acting on synthesis, release or other targets (e.g., receptors), e.g., by interferon β -1b |
| 6.- | oligodendrocyte injury with acute myelin breakdown and degradation of axonal lamellae by microglia/macrophages | prevent myelin injury, e.g., with external competing antigens, such as CoP I |
| 7.- | acute conduction block, with functional impairment | act on ion channels, e.g., potassium channel blockers |
| 8.- | chronic de- and re-myelination process, functional adaptation and reorganization within the brain | enhance oligodendrocyte remyelination mechanisms, e.g., by glial growth factors, or Schwann cell implantation |
| 10 | 9.- persisting lesion without or with astrocyte scar and functional impairment, depending on size and location | symptomatic therapies against spasticity, fatigue, urinary problems, etc., therapeutic aids and training of remaining skills |

- 6 -

While many compounds can attack or another of these MS mechanisms and have, as a rule, typical risks and side effects associated with any specific mechanism, our experimental studies have, surprisingly, demonstrated that inhibitors of phosphodiesterase IV, such a rolipram, are very highly effective on human MS cells, as well as, in different animal models of the disease (e.g., EAE, EAN in different species); furthermore, in different human conditions, their safety and efficacy on different biological parameters has already been observed. With these combined mechanisms it is obvious that low dosages can be used which in monotherapy, or even more so in combination with other maintenance therapies, can be used to prevent relapses or reduce or treat exacerbations of MS with very minor side effects (in contrast to other therapies).

Furthermore, at the same or lower (exceptionally also higher) dosages these drugs can be combined with other compounds as described from 1-9 to achieve additive and synergistic therapeutic effects in order to achieve maximum efficacy (and thus inhibition of progression as described previously) with a minimum of side effects.

An aspect of the present invention thus relates to a method of treating or preventing MS comprising administering an effective amount of a combination of a Type IV phosphodiesterase inhibitor (PDE IV inhibitor) and antiinflammatory or immunomodulatory drugs.

The phosphodiesterase PDE inhibitors suitable for use in this invention are preferably cycloadenosine-3',5'-monophosphate (cAMP) PDE type IV (PDE IV) inhibitors according to the modern classification (J.A. Beavo and D.A. Reifsnyder, Trends Pharmacol. Sci. 11; 150-155, 1990) and include, but are not limited to compounds disclosed in U.S. 4,193,629, WO 92/02220; U.S. 4,186,129; EP 247 725; U.S. 5,064,854; N.A. Saccamono et al., J. Med. Chem. 34, 291-298, 1991; F.J. Vinick et al., J. Med. Chem. 34, 86-89, 1991; J. A. Lowe et al., J. Med. Chem. 34, 624-628, 1991; 1,3-dibutyl-3,7-dihydro-7-(2-oxopropyl)-1H-purin-2,6-dione(denbufylline, BRL 30892);

- 7 -

4-[(3-butoxy-4-methoxyphenyl)methyl]-2-imidazolidinone (RO20-1724);

5,6-diethoxybenzo[b]thiophen-2-carboxylic acid (tibenelast, LY 186655);

5 3-ethyl-1-(3-nitrophenyl)-2,4(1H,3H)-chinazolin-2-one (nitraquazone, TVX 2706);

6-(3,6-dihydro-6-methyl-2-oxo-2H-1,3,4-thiadiazine-5-yl)-1-(3,4-dimethoxybenzoyl)-1,2,3,4-tetrahydro-4,4-dimethylchinoline (EMD 54622);


10 1-ethyl-4-[(1-methylethylidene)hydrazino]-1H-pyrazolo[3,4-b]pyridin-5-carboxylic acid ethylester (etazolate);

N-hydroxy-5,6-dimethoxy-benzo[b]thiophene-2-carboximidamid (Org 30029);

15 2-amino-6-methyl-4-propyl-(1,2,4)triazolo[1,5-a]pyrimidine-5(4H)-one (ICI 63197);

6-[4-(difluoromethoxy)-3-methoxyphenyl]-3(2H)-pyridazinone (zardaverine) pentoxifilline; propentofilline; vinpocetine; and the pharmaceutically acceptable salts thereof.

20 Preferred PDE IV inhibitors are racemic and optically active compounds of formula I



wherein

R¹ is C₁₋₆-alkyl, a heterocyclic ring, or OR⁵; and

R⁵ is C₁₋₆-alkyl, C₃₋₇-cycloalkyl, C₂₋₆-alkenyl,

- 8 -

C₃₋₇-alkinyl, C₃₋₇-cycloalkyl-C₁₋₂-alkyl, aryl, aralkyl, a heterocyclic ring or C₁₋₆-alkyl substituted by one or more halogen atoms, hydroxy, carboxy, C₁₋₄-alkoxy, C₁₋₄-alkoxycarbonyl, or an optionally alkyl substituted amino group;

R² is C₁₋₄-alkyl, C₂₋₄-alkenyl, or C₂₋₄-alkinyl;

R³ is a hydrogen atom, C₁₋₆-alkyl, aryl, aralkyl, or aryl optionally substituted by one or two methyl groups or C₁₋₆-alkanoyl;

R⁴ is a hydrogen atom or C₁₋₆-alkyl;

Y is a direct bond or a CH₂ group;

X is CH₂, CH₂-CH₂, NH, or an oxygen atom; and the pharmaceutically acceptable salts thereof.

Preferred compounds of formula I are those wherein,

R² is methyl;

R³ is a hydrogen atom or C₁₋₆-alkanoyl;

R¹ is OR₅; R₅ is C₁₋₆-alkyl, C₃₋₇-cycloalkyl, or 3-tetrahydrofuranyl;

R⁴ is hydrogen or C₁₋₄-alkyl; and

X is a CH₂ group or oxygen.

Especially preferred compounds of formula I are those wherein R³ is hydrogen.

Specifically exemplified is 4-(3-(cyclopentyloxy)-4-methoxyphenyl)-2-pyrrolidinone (rolipram).

The term "alkyl" as used herein include straight or branched alkyl radicals, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, pentyl, 2-methyl-butyl, 2,2-dimethylpropyl and hexyl.

By the term "alkenyl" as used herein is meant to include, but not limited to vinyl, 1-propenyl, 2-propenyl, 2-propinyl or 3-methyl-2-propenyl.

By the term "cycloalkyl" or "cycloalkyl alkyl" as used herein is meant to include groups of 3-7 carbon atoms, such as cyclopropyl, cyclopropylmethyl, cyclopentyl or cyclohexyl.

- 9 -

By the term "aryl" or "aralkyl" as used herein is meant an aromatic ring or ring system of 6-10 carbon atoms, preferably monocycle, such as phenyl, benzyl, phenethyl or naphthyl.

5 By the term "heterocyclic ring" as used herein is meant a saturated ring of 5 to 6 members having a single oxygen, sulfur or nitrogen atom, such as, but not limited to 2- and 3-tetrahydropyranyl, 2- and 3-tetrahydrofuranyl, pyrrolidino, 2- and 3-pyrrolidyl, piperidinino, 2-, 3- and 4- piperidyl and
10 the corresponding N-alkyl pyrrolidyl and piperidyl rings, wherein the alkyl is of 1-4 carbon atoms. Also encompassed within the scope of this invention are heterocyclic rings having more than one hetero atom such as morpholino, piperazino or N-alkyl piperazino.

15 By the term "halo" as used herein is meant all halogens, i.e., chloro, fluoro, bromo and iodo.

The preparation of the compounds of Formula I can be carried out by the procedure outlined in the above-mentioned patents or by U.S. Patent Nos. 4,153,713; 4,186,129; and
20 5,298,628; WO 86/02268; or EP 0 247 725. Rolipram is 4-[(3-cyclopentyloxy)-4-methoxyphenyl]-2-pyrrolidinone. See, e.g., *Merck Index*, 11th edition, pp. 1312-1313. Rolipram and related compounds can be prepared, e.g., according to U.S. Patent No. 4,193,926.

25 The anti-inflammatory and immunomodulatory drugs suitable for use in this invention include but are not limited to:

1. interferon derivatives, e.g., betaserone, β -interferon, β -interferon muteins;
2. prostane derivatives, e.g., compounds disclosed in
30 PCT/DE93/0013, e.g., iloprost, cicaprost;
3. glucocorticoids, e.g., cortisol, prednisolone, methylprednisolone, dexamethasone;
4. immunosuppressives, e.g., cyclosporine A, FK-506, methoxsalene, thalidomide, sulfasalazine,
35 azathioprine, methotrexate;
5. lipoxxygenase inhibitors, e.g., zileutone, MK-886, WY-50295, SC-45662, SC-41661A, BI-L-357;

- 10 -

6. leukotriene antagonists, e.g., compounds disclosed in DE 4009117; German patent application P 42 42 390.2; WO 9201675; SC-41930; SC-50605; SC-51146; LY 255283 (D.K. Herron et al., FASEB J; 2: Abstr. 4729, 1988); LY 223982 (D.M. Gapinski et al., J. Med. Chem. 33: 2798-2813, 1990); U-75302 and analogs, e.g., described by J. Morris et al., Tetrahedron Lett. 29: 143-146, 1988, C.E. Burgos et al., Tetrahedron Lett. 30: 5081-5084, 1989; B.M. Taylor et al., Prostaglandins 42: 211-224, 1991; compounds disclosed in U.S. 5,019,573; ONO-LB-457 and analogs, e.g., described by K. Kishikawa et al., Adv. Prostagl. Thrombox. Leukotriene Res. 21: 407-410, 1990; M. Konno et al., Adv. Prostagl. Thrombox. Leukotriene Res. 21: 411-414, 1990; WF-11605 and analogs, e.g., disclosed in U.S. 4,963,583; compounds disclosed in WO 9118601, WO 9118879; WO 9118880, and WO 9118883;
7. antiinflammatory substances, e.g., NPC 16570, NPC 17923 described by L. Noronha-Blab. et al., Gastroenterology 102 (Suppl.): A 672, 1992; NPC 15669 and analogs described by R.N. Burch et al., Proc. Nat. Acad. Sci. USA 88: 355-359, 1991; S. Pou et al., Biochem. Pharmacol. 45: 2123-2127, 1993;
8. peptide derivatives, e.g., ACTH and analogs; soluble TNF-receptors; TNF-antibodies; soluble receptors of interleukines, other cytokines, T-cell-proteins; antibodies against receptors of interleukines, other cytokines, T-cell proteins;
9. calcipotriols and their analogues as activators of syntheses of different nerve growth factors, or these growth factors themselves or small peptides thereof which stimulate oligodendrocyte growth (or prevent their apoptosis or destruction) and enhance remyelination.

- 11 -

Our data show that in their effects on human MS cells as well as in different animal models of demyelinating disease (e.g., different EAE models) various of these new drugs can be combined successfully to achieve better protection with less side effects.

By "immunomodulatory drugs", it is meant, e.g., agents which act on the immune system, directly or indirectly, e.g., by stimulating or suppressing a cellular activity of a cell in the immune system, e.g., T-cells, B-cells, macrophages, or other APC cells, or by acting upon components outside the immune system which, in turn, stimulate, suppress, or modulate the immune system, e.g., hormones, receptor agonists or antagonists, and neurotransmitters; immunomodulators can be, e.g., immunosuppressants or immunostimulants.


By "anti-inflammatory drugs", it is meant, e.g., agents which treat inflammatory responses, i.e., a tissue reaction to injury, e.g., agents which treat the immune, vascular or lymphatic systems.

Again, in this new combined therapeutic strategy, there are also many steps where the presumed cascade of MS can be attacked by existing or future drugs (see simplified schedule) after the initial specific or non-specific causes, which include impairment of suppressor T-cell effects on myelin-specific autoimmune T-cells:

The present invention also relates to a method of treating or preventing MS, comprising administering an effective amount of compound of formula I or II alone, preferably Rolipram, a Type IV phosphodiesterase.

The invention, in one aspect, relates to racemates and optically active 4-(polyalkoxyphenyl)-2-pyrrolidones of formula II which are useful in preventing or treating multiple sclerosis:

- 12 -



wherein R₁ and R₂ each are alike or different and are hydrocarbon of up to 18 carbon atoms with at least one being other than methyl, a heterocyclic ring, or alkyl of 1-5 carbon atoms which is substituted by one or more of halogen atoms, hydroxy, carboxy, alkoxy, alkoxycarbonyl or an amino group; R'
5 is a hydrogen atom, alkyl, aryl or acyl; and X is an oxygen atom or a sulfur atom.

These compounds and the methods of making them are described, e.g., in U.S. Patent No. 4,193,926 and WO 92/02220.

A preferred compound according to the present invention is Rolipram. Rolipram is 4-[(3-cyclopentyloxy)-4-methoxyphenyl]-2-pyrrolidinone. See, e.g., *Merck Index*, 11th edition, pages 1312-1313. It is commercially available from
15 Schering AG, Berlin, Germany, or may be prepared, e.g., according to Belgian Patent No. 826,923 or U.S. Patent No. 4,193,926. It is useful conventionally as an antidepressant, e.g., U. Schwabe et al., *Mol. Pharmacol.* 12, 900 (1976); H. Wachtel, *Neuropharmacol.* 22, 267 (1983); H. Wachtel and H.
20 Schneider, *Neuropharmacol.* 25, 1119 (1986); W. Krause and G. Kühne, *Xenobiotica* 18, 561 (1988). Clinical evaluation of Rolipram for depression is reported in E. Zeller et al., *Pharmacopsychiatry* 17, 188 (1984). A comparative clinical trial with amitriptyline, q.v. in severe depressions is
25 reported in F. Eckmann et al., *Curr. Ther. Res.* 43, 291 (1988). Derivatives of Rolipram can also be used according to the invention, i.e., compounds which are structurally related to Rolipram, and are effective in preventing and/or treating MS, e.g., those of formula I.

- 13 -

The present invention also generally relates to the use of a Type IV phosphodiesterase inhibitor, preferably a compound of formula I, especially Rolipram, in multiple sclerosis (MS), for preventing, and/or ameliorating the severity, symptoms, and/or periodicity of recurrence of the disease, e.g., lengthening the time period between episodes in which symptoms flare, and/or suppressing the ongoing immune or autoimmune response associated with the disease.

The invention thus relates to the administration of an effective amount of such a compound, e.g., one according to formula I or II, preferably Rolipram, to a patient to prevent or treat MS. The amount of said compound, e.g., Rolipram, administered is an amount which is effective, for example, in preventing or ameliorating the symptoms of the disease or the disease's recurrence, or affecting the ultimate course of the disease, e.g., blocking the inflammatory response in the brain, the appearance of inflammatory lesions, neuronal or neuroglia cell death, and/ or demyelination and the symptoms typically associated with pathogenesis of the disease.

The present invention also provides pharmaceutical compositions comprising a compound according to formula I or II, preferably a Type IV phosphodiesterase inhibitor, preferably Rolipram, or combinations of drugs as described above, which are useful in preventing or treating multiple sclerosis. According to the method, a compound of formula I or II, or drug combinations, can be administered, e.g., in a single dose, in multiple doses, e.g., through-the-skin injection or by sustained release means such as an implanted osmotic pump.

According to the present invention, a pharmaceutical composition of Formulae I and II, or combinations as described above, comprising an effective amount of each compound described can be administered to patients having multiple sclerosis, e.g., multiple sclerosis variants such as Neuromyelitis Optica (Devic's Disease), Diffuse Sclerosis, Transitional Sclerosis, Acute Disseminated Encephalomyelitis, and Optic Neuritis.

- 14 -

Symptoms of MS which are prevented or ameliorated or treated include: weakness and/or numbness in one or more limbs; tingling of the extremities and tight band-like sensations around the trunk or limbs; dragging or poor control of one or both legs to spastic or ataxic paraparesis; hyperactive tendon reflexes; disappearance of abdominal reflexes; Lhermitte's sign; retrobulbar or optic neuritis; unsteadiness in walking; brain stem symptoms (diplopia, vertigo, vomiting); disorders of micturition; hemiplegia; trigeminal neuralgia; other pain syndromes; nystagmus and ataxia; cerebellar-type ataxia; Charcot's triad; diplopia; bilateral internuclear ophthalmoplegia; myokymia or paralysis of facial muscles; deafness; tinnitus; unformed auditory hallucinations (because of involvement cochlear connections); vertigo and vomiting (vestibular connections); transient facial anesthesia or of trigeminal neuralgia; bladder dysfunction; euphoria; depression; dementia, dull, aching pain in the low back; sharp, burning, poorly localized pains in a limb or both legs and girdle pains; abrupt attacks of neurologic deficit; dysarthria and ataxia; paroxysmal pain and dysesthesia in a limb; flashing lights; paroxysmal itching; and/or tonic seizures, taking the form of flexion (dystonic) spasm of the hand, wrist, and elbow with extension of the lower limb. A patient having MS may have one or more of these symptoms or other clinical manifestations typically associated with MS and one or more can be ameliorated by administering of compounds according to the present invention.

The administration of Type IV phosphodiesterase inhibitors such as Rolipram, or combinations of the latter other drugs, can also block or reduce the physiological and pathogenic deterioration associated with MS, e.g., inflammatory response in the brain and other regions of the nervous system, breakdown or disruption of the blood-brain barrier, appearance of lesions in the brain, tissue destruction, demyelination, autoimmune inflammatory response, acute or chronic inflammatory response, neuronal death, and/or neuroglia death.

- 15 -

The active agents of this invention are useful to treat the different types of MS, including the multifocal, CNS, relapsing and remitting course; the multifocal, CNS, progressive course; the single-site, relapsing and remitting course; and other variants of multiple sclerosis. See, e.g., *Cecil's Textbook of Medicine*, edited by James B. Wyngaarden, 1988.

Effects of the administration of Rolipram and other Type IV phosphodiesterase inhibitors, and combinations of it with other drugs include, e.g., preventing the disease, ameliorating symptoms of the disease, reducing the annual exacerbation rate (i.e., reducing the number of episodes per year), slowing the progression of the disease, or reducing the appearance of brain lesions (e.g., as identified by MRI scan). The episodic recurrence of the mentioned diseases such as MS can be ameliorated, e.g., by decreasing the severity of the symptoms (such as the symptoms described above) associated with the, e.g., MS episode, or by lengthening the time period between the occurrence of episodes, e.g., by days, weeks, months, or years, where the episodes can be characterized by the flare-up and exacerbation of disease symptoms, or preventing or slowing the appearance of brain inflammatory lesions. See, e.g., Adams, R.D., *Principles of Neurology*, 1993, page 777, for a description of a neurological inflammatory lesion.

Other specific, suitable, non-limiting examples of Type IV phosphodiesterase inhibitors which can be employed in this invention include compounds described in WO93/19068, compounds RO 20-1724 (4-[(3-butyloxy-4-methoxyphenyl)methyl]-2-imidazolidinone), ICI 63197 (2-amino-6-methyl-4-propyl[1,2,4]triazolo[1,5-a]pyrimidin-5(4H)-one), denbufylline and etazolate.

By "Type IV phosphodiesterase inhibitor", "specific Type IV phosphodiesterase inhibitor", and similar expressions are meant a selective, i.e., specific, such inhibitor, where the compound binds to or inhibits preferentially the Type IV phosphodiesterase when compared to known types of

- 16 -

phosphodiesterase types, e.g., I, II, or III, e.g., whereby the compound has a lower IC_{50} (more potent) for the Type IV phosphodiesterase, such as where the IC_{50} is, e.g., 2-fold, 5-fold, 10-fold, 50-fold, or more potent, for the Type IV phosphodiesterase compared to another known type of phosphodiesterase, e.g., I, II, or III. Such selectivity of a compound according to the present invention for a Type IV phosphodiesterase can also be conferred by other means, such as the manner in which it is delivered to its target, e.g., the compound can be associated with an agent which targets it to a specific tissue or cell type having the Type IV phosphodiesterase; the manner in which it interacts with the host's metabolism and/or physiology; or synthesizing PDE inhibitor prodrugs where activation of the PDE inhibitor is accomplished by enzymes present in the desired cells or tissues but absent in others.

The specific inhibition of a Type IV phosphodiesterase can be measured conventionally, e.g., according to the methods described in Reeves et al., *Biochem. J.*, 241:535-541, 1977; by macrophage assay, as described, e.g., in Schade et al., *Europ. J. Pharmacol.*, 230:9-14, 1993; or WO 93/19068. For a review of phosphodiesterase specificity and how to determine it, see, e.g., Nicholson et al., *Trends Pharmacol. Sci.*, 12:19-27 (1991).

The activity of this invention of Type IV phosphodiesterase inhibitors such as Rolipram can be detected, for example, in animals suffering from Experimental Allergic Encephalomyelitis (EAE), an experimental T-lymphocyte initiated disease of the CNS. It can be produced, e.g., in rodents, guinea pigs, rabbits, and primates, by, e.g., immunizing animals with myelin, e.g., from a human brain, and/or corticosteroid administration over a long period of time. It can also be produced by injecting an animal with T-lymphocytes obtained from an animal suffering from EAE.

In particular, the activity can be detected in *Callithrix jacchus* (common marmoset) which has been immunized with myelin, e.g., from a human brain. The *Callithrix jacchus*

- 17 -

develops EAE with essentially similar histopathology and neurological symptoms as those at certain stages of the human disease, MS.

The present invention generally relates to the treatment of MS with a combination of a PDE IV inhibitor with an interferon derivative, a prostane derivative, a glucocorticoid, an immunosuppressant, a lipoxxygenase inhibitor, a leukotriene antagonist, an antiinflammatory substance, a peptide derivative or a calcipotriol or analog thereof.

A preferred combination consists of a PDE IV inhibitor and an interferon derivative, a prostane derivate or a leukotriene antagonist, e.g., betaserone, iloprost, cicaprost, or 5-[(E)-(2S)-2-((1E,3E)-(5S)-5-hydroxy-6,6-trimethylene-9-phenyl-1,3-nonadiene-8-ynyl)-cyclohexylidene]-pentanoic acid or esters thereof.

The pharmaceutical compositions according to the present invention are prepared conventionally, comprising substances which are customarily used in pharmaceutical, e.g., see *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing Company (1990), including excipients, carriers, adjuvants and buffers. The compositions can be administered, e.g., parenterally, enterally, orally, intramuscularly, topically, subcutaneously, intravenously, by aerosol, intrathecally directly into the cerebral spinal fluid of the CNS, or preferably by sustained release using, e.g., an implanted mini-osmotic pump (e.g., the ALZET pump manufactured by ALZA Corporation, P. O. Box 10950, Palo Alto, CA 94303), or other routes useful to achieve an effect.

Conventional excipients include pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, enteral or topical application which do not deleteriously react with the agents. Suitable pharmaceutically acceptable adjuvants include, but are not limited to, water, salt solutions, alcohols, gum arabic, vegetable oils, polyethylene glycols, gelatine, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil,

- 18 -

fatty acid monoglycerides and diglycerides, pentaerythritol fatty acid esters, hydroxy-methylcellulose, polyvinyl pyrrolidone, cyclodextrins, etc. The pharmaceutical preparations can be sterilized and, if desired, mixed with stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, flavoring and/or aromatic substances, etc., which do not react deleteriously with the active compounds.

For parenteral application, particularly suitable are injectable sterile solutions, preferably oil or aqueous solutions, as well as suspensions, emulsions or implants, including suppositories. Ampoules are convenient unit dosages.

For enteral application, particularly suitable are tablets, dragees, suppositories or capsules having talc and/or a carbohydrate carrier or binder. The carrier may be lactose, corn starch, potato starch or a combination thereof. A syrup or elixir may be used when a sweetened vehicle is employed.

The compositions can also be formulated in an aqueous solution, optionally with the addition of additives customary in galenicals, for example, buffers; electrolytes such as sodium chloride; antioxidants such as ascorbic acid; adjuvants, e.g., methyl cellulose, lactose and mannitol and/or surfactants, e.g., lecithins and Tweens and/or aromatic substances for flavoring, e.g., ethereal oils.

Amounts of Type IV phosphodiesterase inhibitors and drug combinations can be determined routinely based on the information given herein, e.g., using the EAE model. However, any amount which is effective in treating MS can be administered to ameliorate or treat the disease. Dosages are determined conventionally, see, e.g., *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing Company (1990). The composition may be administered in a single dose unit or in multiple dosages administered, e.g., twice, three, or four times a day, or by an osmotic pump, which delivers the drug(s) continuously. A Type IV phosphodiesterase inhibitor can be administered at the same time as the anti-inflammatory,

- 19 -

immunomodulatory, etc., drug in a single or separate dosage unit, or the drugs can be administered at a different time or, e.g., sequentially.

The exact dose of any component or combination to be administered is determined by the attending clinician and is dependent, e.g., on the potency of the compound administered, the age, weight, condition, and response of the patient.

Generally, PDE IV inhibitors are administered alone in amounts of about 0.005 - 2 mg/kg/day, preferably 0.1 -.7 mg/kg/day or 0.5 mg/kg/day, more preferably 0.005 - 0.1 mg/kg/day and the immuno-modulatory or anti-inflammatory, etc., is administered alone in amounts of, e.g., about 0.01 μ g/kg/day for a prostacyclin or to about 10 mg/kg/day for a steroid. According to the present invention, the latter can be administered in lower doses than would be expected for purely additive effects, e.g., about 0.0005 to about 0.01 mg/kg/day for a PDE IV inhibitor and about 0.001 μ g/kg/day to about 1 mg/kg/day for an immunomodulatory or anti-inflammatory drug.

Since the present invention relates to treatment of MS with a combination of active ingredients wherein said ingredients can be administered separately, the invention also relates to combining separate pharmaceutical composition in kit form. The kit form is particularly advantageous when the separate components are administered in different dosage forms (i.e., oral and parenteral) or are administered at different dosage intervals.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative and not limitative of the remainder of the disclosure in any way whatsoever.

In the foregoing and in the following examples, all temperatures are set forth uncorrected in degrees Celsius; and, unless otherwise indicated, all parts and percentages are by weight.

- 20 -

Brief Description of the Drawings

Various other objects, features, and attendant advantages of the present invention will be more fully appreciated as the same becomes better understood when considered in conjunction with the accompanying drawings, in which like reference characters designate the same or similar parts throughout the several views and wherein:

Figure 1 shows the prevention of Experimental Allergic Encephalomyelitis (EAE) by Rolipram in a marmoset. A, B, and C received Rolipram (10 mg/kg) in DMSO; D and E received an equivalent volume of DMSO. Marmosets immunized with human spinal cord homogenate received either Rolipram or placebo five days after immunization.

Figure 2 shows the treatment with Rolipram of marmoset having EAE.

Examples

EXAMPLE 1

Rolipram was produced by Schering AG (Berlin) and is comprised of (+) and (-) racemates of 4-[(3-cyclopentyloxy)-4-methoxyphenyl]-2-pyrrolidinone. It was dissolved in dimethylsulfoxide (DMSO) at 20 mg/ml.

Human brain white matter homogenate was prepared from autopsy material in complete Freund's adjuvant (CFA) containing *M. tuberculosis* (strain H37 Ra).

Bordetella pertussis vaccine was obtained from the Massachusetts Public Health Department, Biological Laboratories, Boston, Massachusetts.

Zofran was obtained from the University of California Medical Center Pharmacy.

The marmosets were purchased from the New England Regional Primate Research Center and were maintained and cared for in accordance with the guidelines of the Internal Animal Care and Use Committee of the University of California, San Francisco.

Animals were immunized with human brain white matter homogenate (200 mg) in CFA containing 3 mg/ml killed *M.*

- 21 -

tuberculosis (H37 Ra strain) by intradermal injection (0.6 ml) over four sites on the dorsal axilla and inguinal region. On the day of immunization and again 2 days later 10×10^{10} inactivated Bordetella pertussis (Bordetella pertussis vaccine) were infused intravenously in 10 ml saline.

On day 5, following immunization, animals were injected subcutaneously in the back of the neck with DMSO (placebo) or with DMSO containing Rolipram to give a dose of 10 mg/kg. Treatment with DMSO or DMSO with Rolipram was preceded 20 min. by an injection of 0.3-0.6 mg/kg Zofran intramuscularly (Odansetron Hydrochloride, Glaxo) to prevent salivation, vomiting, excessive grooming, and head twists. Such treatments were repeated every 48 hours throughout the study.

Animals were observed daily and subjected to a standardized scoring system to record the severity of clinical symptoms:

0. Normal
1. Lethargy, anorexia, weight loss
2. Ataxia, tremor
3. Blindness, paraplegia or hemiplegia
4. Quadraparesis or quadriplegia
5. Moribund

At various times, animals were anesthetized and subjected to MRI.

Experiment 1: Prevention of EAE by Rolipram Treatment

Marmosets were immunized with spinal cord homogenate as described. On day 5, following immunization, three marmosets received Rolipram (10 mg/kg) in DMSO. See Figure 1, A, B, and C. Two marmosets received an equivalent volume of DMSO after the same interval. See Figure 1, D and E. The treatment was repeated every 48 hours.

The animals treated with DMSO (placebo) developed clinical symptoms consistent with EAE 15 days following immunization; see Figure 1, D and E. None of the Rolipram treated animals developed symptoms during the 8 week interval of observation; see Figure 1, A, B, and C.

- 22 -

Magnetic Resonance Imaging (MRI) analysis showed that the two animals with EAE symptoms developed one or more lesions in the brain which "enhanced" with Magnevist (gadolinium, DTPA), indicating an active edematous response consistent with the vascular inflammatory lesions seen in EAE or MS (Alvord, etc.). None of the Rolipram treated animals developed detectable lesions during the 8 week interval of observation.

Development of EAE After Withdrawal of Rolipram

On day 60, following immunization, the treated animals were removed from treatment and observed for signs of EAE. Two marmosets began to show clinical signs of EAE on day 17, following withdrawal of Rolipram.

Experiment 2: Treatment of Active EAE with Rolipram

In the previous experiment, it was clearly demonstrated that Rolipram could prevent EAE. It is of interest to determine if Rolipram can also affect active EAE. In this experiment, a marmoset was immunized as previously described, allowed to develop symptoms of chronic EAE and subsequently treated with Rolipram. The animal was treated with escalating doses of Rolipram in DMSO administered as described in the previous experiment. The physical symptoms were monitored as described and MRI analysis was done at times before and after treating (Figure 2).

The animal showed marked improvement on day 10 after initiating treatment. The animal showed MRI improvement on day 14 following initiating Rolipram treatment and, from that time, the condition stabilized with slower improvements.

The results of Experiment 1 indicate that Rolipram treatment blocked the neurological signs of EAE. The MRI results indicate that the inflammatory response was blocked and demyelination did not occur. The untreated control animals developed clear signs of EAE and inflammatory lesions as indicated by MRI analysis. The fact that treated animals developed EAE when removed from treatment showed that the immune response to brain homogenate had occurred sufficiently

- 23 -

to initiate the disease; however, some subsequent step in pathogenesis was blocked.

The results of Experiment 2 indicate that Rolipram treatment can inhibit active disease.

5 EXAMPLE 2

To assess the ability of rolipram alone or in various combinations as described to modify autoimmune processes, we investigated its influence on TNF production *in vitro* by MBP-specific T-cell lines from MS patients and Lewis rats. MBP is
10 a major candidate antigen in MS, and T-cell-mediated immunity is of crucial importance in its pathogenesis. Similar to EAE, MBP-specific T-cells in humans are often cytotoxic, of Th1 type secreting interferon (IFN)-gamma and TNF/LT, and recognize epitopes that are also encephalitogenic in EAE.

15 Rolipram selectively inhibited TNF production by human MBP-specific T-cell lines (TCL) in a dose-dependent manner alone and in combination.

Similar results were found using an encephalitogenic CD4+ MBP-specific rat TCL (L1402). TNF/LT (lymphotoxin) production
20 measured in a cytotoxicity bioassay was inhibited in a dose range comparable to the human lines. Moreover, inhibition was stereospecific, with the (-)-enantiomer being 55 times more effective than the (+)-enantiomer. The EC₅₀ of (-)-rolipram, (+)-rolipram, and (-)-rolipram given alone were 20 nM, 280 nM,
25 and 1100 nM, respectively. Previous investigations had shown that inhibition of cAMP PDE by rolipram is stereospecific. *In vitro* and *in vivo* binding data in mouse and rat forebrain tissue with ³H-rolipram proved for the (-)-enantiomers a 15-30
30 times higher affinity than the (+)-enantiomers. In line with these findings, our data strongly suggest that rolipram inhibits TNF/LT production in human and rat autoreactive T-cells by an intracellular cAMP PDE dependent mechanism.

TNF and LT may both be produced by autoactive T-cells. CD4+ cells have been reported to be the major source of TFN in
35 autoimmune insulitis of NOD mice. The cytokine bioassay for

- 24 -

TNF/LT detection employed here is sensitive to TNF and LT, but is 200 times more sensitive to the former.

The results of our *in vivo* findings prompted us to perform treatment experiments in EAE after active immunization (aEAE) and adoptive cell transfer (tEAE) in Lewis rats. When rolipram was administered from day 7 through day 23 in aEAE as monotherapy or in various combinations, the appearance of neurological symptoms was completely prevented. In clinically manifest EAE, treatment was started within 6 hours of the onset of symptoms. In the treated group, disease did only progress moderately, whereas the controls developed severe EAE. None of the rolipram-treated animals developed grade 4 (paraplegia), whereas 4 to 7 animals in the vehicle-treated control group reached this level of impairment in one typical experiment. In tEAE similar effects were observed.

Prophylactic treatment resulted in only minor symptoms with a mean maximum score (MMN) of 0.3 ± 0.11 ($n = 5$) in the treated group, as compared to 2.5 ± 0.25 ($n = 5$) in the controls ($p < 0.01$). Treatment after onset of symptoms also lead to a marked reduction in maximum severity (treated group: MMS 0.7 ± 0.10 , $n = 5$; vehicle treated matched controls 2.45 ± 0.56 , $n = 5$, $p < 0.01$).

In order to distinguish further between a long-term prophylactic or a temporary suppressive effect, animals received the drug from the day of active immunization until day 11. With this regimen aEAE in treated animals was delayed by 3.5 days, but disease severity and duration was otherwise similar compared to the controls. This indicates that suppression of EAE and presumably TNF/LT production is temporary, and deletion of autoreactive T-cells by rolipram, as, e.g., in the case of cyclophosphamide, is therefore unlikely.

Histological analysis was performed on selected animals with aEAE. There were only few and mild cellular lesions in the prophylactically treated animals. By contrast, two or three animals that were treated after onset of clinical signs showed cellular infiltrates similar to those in the controls.

- 25 -

Previous investigations have shown that inflammatory infiltrates in the central nervous system do not necessarily correlate with the degree of neurological deficit. In a study on EAE, induced by myelin oligodendrocyte glycoprotein (MOG)-specific T-cells, the lack of neurological signs was ascribed to a decrease of macrophages and parenchymal inflammation, whereas perivascular inflammation and the synthesis of TNF, IFN-gamma, and interleukin-6 was clearly present. In our system, however, the timely onset of paralysis and the morphologically similar appearance of infiltrates in some of the treated animals argues against such a phenomenon. We propose that during rolipram treatment of clinically manifest EAE, suppression of local TNF production in addition to its other effects is crucial regardless of the discrepancy between the histological and clinical scores. Using in-situ hybridization, it was shown recently that the present of TNF expressing cells in the CNS correlates well with the clinical signs in EAE. Using this approach it should be possible to identify the rolipram-sensitive cell types in EAE and MS lesions - inflammatory cells, glial cells, or both.

In contrast, rolipram by its many effects is expected to be of great therapeutic use in MS and other patients, especially in appropriate combinations, as shown by these and other experiments and as indicated by the considerations already discussed.

The preceding examples can be repeated with similar success by substituting the generically or specifically described reactants and/or operating conditions of this invention for those used in the preceding examples.


- 26 -

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention and, without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

5

CLAIMSWe claim:

1. A pharmaceutical composition for treating MS, Guillan-Barre's Syndrome, virus-, bacteria-, or parasite-related demyelinating diseases, encephalopathies related to HIV, meningococcal or toxoplasma infections, central malaria, or Lyme's Disease, comprising an effective amount of a combination of a PDE IV inhibitor and an anti-inflammatory or immunomodulatory drug and a pharmaceutically acceptable carrier.
2. A pharmaceutical composition according to claim 1 for treating MS.
3. A composition of claim 1, wherein the PDE IV inhibitor is a compound according to formula I



wherein

R^1 is C_{1-6} -alkyl, a heterocyclic ring, or OR^5 ; and

- 28 -

- R⁵ is C₁₋₆-alkyl, C₃₋₇-cycloalkyl, C₂₋₆-alkenyl, C₃₋₇-alkinyl, C₃₋₇-cycloalkyl-C₁₋₂-alkyl, aryl, aralkyl, a heterocyclic ring or C₁₋₆-alkyl substituted by one or more halogen atoms, hydroxy, carboxy, C₁₋₄-alkoxy, C₁₋₄-alkoxycarbonyl, or an optionally alkyl substituted amino group;
- R² is C₁₋₄-alkyl, C₂₋₄-alkenyl, or C₂₋₄-alkinyl;
- R³ is a hydrogen atom, C₁₋₆-alkyl, aryl, aralkyl, or aryl optionally substituted by one or two methyl groups or C₁₋₆-alkanoyl;
- R⁴ is a hydrogen atom or C₁₋₆-alkyl;
- Y is a direct bond or a CH₂ group;
- X is CH₂, CH₂-CH₂, NH, or an oxygen atom; and pharmaceutically acceptable salts thereof.

4. A composition of claim 1, wherein the PDE IV inhibitor is Rolipram.

5. A composition of claim 1, wherein the anti-inflammatory or immunomodulatory drug is an interferon derivative, a prostane compound, a leukotriene antagonist, a peptide compound, or a calcipotriol or analog thereof.

6. A composition of claim 1, wherein the anti-inflammatory or immunomodulatory drug is Iloprost.

7. The use of a PDE IV inhibitor in combination with a anti-inflammatory or immunomodulatory drug for the preparation of a pharmaceutical composition useful in the treatment of MS.

8. A method of preventing or treating multiple sclerosis comprising administering to a host in need thereof an effective amount of a composition according to claim 1.


9. A method according to claim 8, wherein the time between or the severity of symptoms of the episodic recurrences of the multiple sclerosis is ameliorated.

- 29 -

10. A method according to claim 8, wherein an inflammatory lesion associated with said multiple sclerosis is prevented or treated.

11. A method according to claim 8, wherein the appearance of an inflammatory lesion associated with said multiple sclerosis is slowed.

12. A method of preventing or treating multiple sclerosis, comprising administering to a host in need thereof an effective amount of a compound according to formula II



wherein:

R₁ and R₂ each are alike or different and are C₁₋₁₈-alkyl with at least one being other than methyl, a heterocyclic ring, or C₁₋₅-alkyl substituted by one or more of halogen atoms, hydroxy, carboxy, alkoxy, alkoxycarbonyl or an amino group;

R' is a hydrogen atom, alkyl, aryl or acyl; and X is an oxygen atom or a sulfur atom.

13. A method of claim 12, wherein said compound is 4-[(3-cyclopentyloxy)-4-methoxyphenyl]-2-pyrrolidinone.

14. A method according to claim 12, wherein one of R₁ and R₂ is methyl, and the other is C₃₋₇-cycloalkyl.

- 30 -

15. A method of claim 12, wherein the time between or the severity of symptoms of episodic recurrences of multiple sclerosis is ameliorated.

16. A method according to claim 12, wherein an inflammatory lesion associated with said multiple sclerosis is prevented or treated.

17. A method according to claim 12, wherein the appearance of an inflammatory lesion associated with said multiple sclerosis is slowed.

18. A method according to claim 12, wherein said compound is a Type IV phosphodiesterase.

19. A method of preventing or treating multiple sclerosis comprising administering to a host in need thereof an effective amount of a Type IV phosphodiesterase inhibitor.

20. A method according to claim 19, wherein the time between or the severity of symptoms of the episodic recurrences of the multiple sclerosis is ameliorated.

21. A method according to claim 19, wherein an inflammatory lesion associated with said multiple sclerosis is prevented or treated.

22. A method according to claim 19, wherein the appearance of an inflammatory lesion associated with said multiple sclerosis is slowed.

23. A method of preventing or treating multiple sclerosis comprising administering to a host in need thereof an effective amount of 4-[(3-cyclopentyloxy)-4-methoxyphenyl]-2-pyrrolidinone.

- 31 -

24. A kit comprising in separate containers in a single package pharmaceutical compositions for use in combination to treat MS which comprises a PDE IV inhibitor and, in a second container, a pharmaceutical composition comprising an anti-inflammatory and immunomodulatory drug.

1/2




FIG. 1




FIG. 2

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 95/00332

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/40 //(A61K31/40, 31:00, 38:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	J. NEUROBIOLOGY, vol. 54, no. 1-2, 1994 page 198 SOMMER ET AL. 'Tumor-necrosis factor-directed therapy of autoimmune encephalomyelitis by the phosphodiesterase IV inhibitor rolipram' * see abstract no. P15.08 ---	12-23
X	ACTA NEUROLOGICA SCANDINAVICA, vol. 88, no. 2, 1993 pages 97-99, NATAF ET AL. 'Pentoxifylline inhibits experimental allergic encephalomyelitis' * see whole document *	19-22
Y	* " *	7-23
	--- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

21 July 1995

Date of mailing of the international search report

21.08.95.

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Isert, B

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IB 95/00332

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EUR. J. PHARMACOL., vol. 230, no. 1, 5 January 1993 pages 9-14, SCHADE ET AL. 'The specific Type III and IV Phosphodiesterase inhibitor zardaverine suppresses formation of tumor necrosis factor by macrophages' * see whole document, in particular Fig. 2 and Table 2*	1-6,24
Y	* " *	7-23
Y	WO,A,92 19594 (SMITH-KLINE BEECHAM) 12 November 1992 * see claims 1-12 * -----	12-23

INTERNATIONAL SEARCH REPORT

Information on patent family members

Information on patent family members

Internal Application No

PCT/IB 95/00332

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9219594	12-11-92	AU-A- 1917092	21-12-92
		CN-A- 1067244	23-12-92
		EP-A- 0584208	02-03-94
		JP-T- 6507405	25-08-94

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 213/30, A61K 31/44	A1	(11) International Publication Number: WO 95/35282 (43) International Publication Date: 28 December 1995 (28.12.95)
(21) International Application Number: PCT/GB95/01460 (22) International Filing Date: 21 June 1995 (21.06.95) (30) Priority Data: 9412383.3 21 June 1994 (21.06.94) GB (71) Applicant: CELLTECH THERAPEUTICS LIMITED [GB/GB]; 216 Bath Road, Slough, Berkshire SL1 4EN (GB). (72) Inventors: HEAD, John, Clifford; Flat 2A, 8 Claremont Road, Windsor, Berkshire SL4 3AX (GB). WARRELOW, Graham, John; Oakside, 4 Wieland Road, Northwood, Middlesex HA6 3QU (GB). ALEXANDER, Rikki, Peter; 14 Carrington Road, High Wycombe, Bucks HP12 3HY (GB). BOYD, Ewan, Campbell; 30 Ledi Avenue, Tullibody, Clacks FK10 2RZ (GB). (74) Agent: SKAILES, Humphrey, John; Frank B. Dehn & Co., Imperial House, 15-19 Kingsway, London WC2B 6UZ (GB).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i>
(54) Title: (R)-4-(2-(3-CYCLOPENTYLOXY-4-METHOXYPHENYL)-2-PHENYLETHYL) PYRIDINE HYDROGEN SULPHATE SALT AS PDE TYPE IV INHIBITOR		
(57) Abstract		
<p>The hydrogen sulphate salt of the selective phosphodiesterase type IV inhibitor (R)-4-[2-(3-Cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]-pyridine hydrogen sulphate salt is described. The salt possesses a number of advantageous chemical and physical characteristics making it particularly suitable for pharmaceutical formulation for use in <i>inter alia</i> the prophylaxis and treatment of asthma.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

(R)-4-(2-(3-CYCLOPENTYLOXY-4-METHOXYPHENYL)-2-PHENYLETHYL) PYRIDINE HYDROGEN SULPHATE SALT AS PDE TYPE IV INHIBITOR.

5 This invention concerns (R)-4-[2-(3-cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine hydrogen sulphate salt, to processes for its preparation, to pharmaceutical compositions containing it and to its use in medicine.

10 In our International Patent Specification No. WO 94/14742 we describe a series of tri-substituted phenyl derivatives which are potent inhibitors of the phosphodiesterase (PDE) type IV isoenzyme at concentrations at which they have little or no inhibitory action on other PDE isoenzymes. The compounds are of use in medicine, especially in the prophylaxis and
15 treatment of asthma.

A particularly useful member of the series is (R)-4-[2-(3-cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine, hereinafter also referred to as CT1730. In WO 94/14742 we describe the production of CT1730 as the
20 free base [Example 16 (i)] and generally disclose salts of the compound. The production of the racemate is also described [Example 3a)], together with the production of the corresponding hydrochloride salt.

For the use in medicine of a compound such as CT1730 it is essential that
25 a form is available which has appropriate chemical and physical characteristics which enable the easy preparation of stable pharmaceutical formulations. The free base form of CT1730 is unsuitable for this purpose, but we have now found a particular salt form of the compound which has advantageous chemical and physical characteristics. The salt is
30 particularly suitable for pharmaceutical formulation, especially when compared to other salt forms of CT1730.

Thus according to one aspect of the invention we provide (R)-4-[2-(3-cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine hydrogen
35 sulphate salt.

The hydrogen sulphate salt of the invention possesses a number of advantageous chemical and physical characteristics. In particular, it is (1) highly crystalline; (2) thermally stable; (3) non-hygroscopic, and (4) soluble in aqueous solutions over a wide range of concentrations. In addition, it is readily prepared free of solvent and other impurities.

All of these properties provide the salt of the invention with easy preparation, handling, purity and stability characteristics which make it a particularly suitable form for pharmaceutical formulation. In contrast, other salt forms of CT1730, particularly the hydrochloride, hydrobromide, hydroiodide, methanesulphonate, p-toluenesulphonate, besylate, phosphate, sulphate, nitrate, maleate and fumarate, lack one or more of these properties and are unsuitable for formulation.

The salt according to the invention may be prepared from the corresponding CT1730 free base or a salt thereof. Thus, according to a further aspect of the invention, we provide a process for the preparation of (R)-4-[2-(3-cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine hydrogen sulphate salt which comprises reacting (R)-4-[2-(3-cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine or a salt thereof with sulphuric acid.

The reaction may be performed in any suitable solvent or mixtures of solvents. Particular solvents include alcohols, such as ethanol or isopropyl alcohol, and aromatic hydrocarbons such as benzene and toluene. In general, the reaction may be performed at around ambient temperature or above, for example up to around 50°C.

In this reaction, the starting material is preferably the free base. However another salt may be used, or if desired a mixture of the free base and the other salt.

The free base or salt starting materials for this reaction may be prepared by the methods described in International Patent Specification No. WO 94.14742, or in our International Patent Application No. PCT/GB

94/02799. In these applications the free base is described as above or as (+)-4-[2-(3-cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine.

5 The salt according to the invention is a selective and potent inhibitor of PDE IV. The ability of the compound to act in this way may be simply determined by the tests described in the Examples hereinafter.

10 The salt according to the invention is thus of particular use in the prophylaxis and treatment of human diseases where an unwanted inflammatory response or muscular spasm (for example bladder or alimentary smooth muscle spasm) is present and where the elevation of cAMP levels may be expected to prevent or alleviate the inflammation and relax muscle.

15 Particular uses to which the salt of the invention may be put include the prophylaxis and treatment of asthma, especially inflamed lung associated with asthma, cystic fibrosis, or in the treatment of inflammatory airway disease, chronic bronchitis, eosinophilic granuloma, psoriasis and other benign and malignant proliferative skin diseases, endotoxic shock, septic
20 shock, ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, inflammatory arthritis, chronic glomerulonephritis, atopic dermatitis, urticaria, adult respiratory distress syndrome, diabetes insipidus, allergic rhinitis, allergic conjunctivitis, vernal conjunctivitis, arterial restenosis and arteriosclerosis.

25 The salt of the invention also suppresses neurogenic inflammation through elevation of cAMP in sensory neurones. It is, therefore, analgesic, anti-tussive and anti-hyperalgesic in inflammatory diseases associated with irritation and pain.

30 The salt according to the invention may also elevate cAMP in lymphocytes and thereby suppress unwanted lymphocyte activation in immune-based diseases such as rheumatoid arthritis, ankylosing spondylitis, transplant rejection and graft versus host disease.

35

The salt according to the invention have also been found to reduce gastric acid secretion and therefore can be used to treat conditions associated with hypersecretion.

- 5 The salt of the invention suppresses cytokine synthesis by inflammatory cells in response to immune or infectious stimulation. It is, therefore, useful in the treatment of bacterial, fungal or viral induced sepsis and septic shock in which cytokines such as tumour necrosis factor (TNF) are key mediators. Also the salt of the invention suppresses inflammation and
10 pyrexia due to cytokines and is, therefore, useful in the treatment of inflammation and cytokine-mediated chronic tissue degeneration which occurs in diseases such as rheumatoid or osteo-arthritis.

- Over-production of cytokines such as TNF in bacterial, fungal or viral
15 infections or in diseases such as cancer, leads to cachexia and muscle wasting. The salt of the invention ameliorates these symptoms with a consequent enhancement of quality of life.

- The salt of the invention also elevates cAMP in certain areas of the brain
20 and thereby counteract depression and memory impairment.

- The salt of the invention suppresses cell proliferation in certain tumour cells and can be used, therefore, to prevent tumour growth and invasion of normal tissues.

- 25 For the prophylaxis or treatment of disease the salt according to the invention is particularly suitable for formulation as a pharmaceutical composition. Thus according to a further aspect of the invention we provide a pharmaceutical composition which comprises (R)-4-[2-(3-cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine hydrogen
30 sulphate salt together with one or more pharmaceutically acceptable carriers, excipients or diluents.

- Pharmaceutical compositions according to the invention may take a form
35- suitable for oral, buccal, parenteral, nasal, topical or rectal administration, or a form suitable for administration by inhalation or insufflation.

Advantageously, the compositions may be prepared using conventional procedures and thus the invention further provides a process for the preparation of a pharmaceutical composition containing (R)-4-[2-(3-cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine hydrogen sulphate salt together with one or more pharmaceutically acceptable carriers, excipients or diluents which comprises the step or steps of bringing into contact (R)-4-[2-(3-cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine hydrogen sulphate salt with one or more pharmaceutically acceptable carriers, excipients or diluents

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets, lozenges or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium glycollate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents, emulsifying agents, non-aqueous vehicles and preservatives. The preparations may also contain buffer salts, flavouring, colouring and sweetening agents as appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

The salt according to the invention may be formulated for parenteral administration by injection e.g. by bolus injection or infusion. Formulations

for injection may be presented in unit dosage form, e.g. in glass ampoule or multi dose containers, e.g. glass vials. The compositions for injection may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as
5 suspending, stabilising, preserving and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

10 In addition to the formulations described above, the salt according to the invention may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation or by intramuscular injection.

15 For nasal administration or administration by inhalation, the salt according to the present invention is conveniently delivered in the form of an aerosol spray presentation for pressurised packs or a nebuliser, with the use of suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas or mixture of gases.

20 The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack or dispensing device may be accompanied by instructions for administration.

25 The quantity of the salt of the invention required for the prophylaxis or treatment of a particular inflammatory condition will vary depending on the condition of the patient to be treated. In general, however, daily dosages may range from around 100ng/kg to 100mg/kg, e.g. around 0.01mg/kg to
30 40mg/kg body weight for oral or buccal administration, from around 10ng/kg to 50mg/kg body weight for parenteral administration and around 0.05mg to around 1000mg e.g. around 0.5mg to around 1000mg for nasal administration or administration by inhalation or insufflation.

35 The following Examples illustrate the invention.

EXAMPLE 1**(R)-4-[2-(3-Cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine
hydrogen sulphate salt**

(R)-4-[2-(3-Cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine

- 5 (14.5g, 39mmol; prepared as described in International Patent Application
No. PCT/GB94/02799 was dissolved in warm ethanol (150ml) and the
clear solution then cooled to room temperature. Concentrated sulphuric
acid (3.3ml, 60mmol) was added with swirling over one minute, followed
by a few seed crystals. The solution was allowed to stand at room
10 temperature for 1.5 hours during which time needle-like crystals steadily
developed. The solution was then left at 40°C overnight to maximise
yield. The resulting product was warmed to room temperature and the
crystalline product was collected by suction filtration with *t*-butylmethyl
ether washing. Once sufficiently dry, the product was transferred to a
15 vacuum oven and heated *in vacuo* to dryness (65°C, ~0.05mbar,
overnight) to afford the title salt as a white crystalline powder (16.2g); m.p.
144-146°C; Found C, 63.72; H, 6.15; N, 2.97. C₂₅H₂₉NO₆S requires C,
63.68; H, 6.20; N, 2.97%.

- 20 The title salt (3.3g) was recrystallised (with slow cooling to room
temperature, then leaving at room temperature for 2 hours) from absolute
ethanol (~40ml). The resulting white needles were filtered, washed with
diethyl ether and dried at 75°C at 0.05mbar overnight.

25 EXAMPLE 2

This example shows the formulation of granules containing (R)-4-[2-
(3-cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine hydrogen
sulphate salt (referred to in this example as 'Active Ingredient'):

30

	<u>Unit Quantity</u>	<u>Batch Quantity</u>
Active Ingredient	0.1263mg	0.2210g
Maize Starch BP	112.7mg	197.2g
Purified Water	-	110ml
35 Colloidal silica (Aerosil 200)	1.71mg	2.993g

The granules were used to either fill capsules, or were compressed into tablets.

5 **EXAMPLE 3**

The activity and selectivity of the salt according to the invention was demonstrated in the following tests. In these tests the abbreviation FMLP represents the peptide N-formyl-met-leu-phe.

10 **Isolated Enzyme**

The potency and selectivity of the salt of the invention was determined using distinct PDE isoenzymes as follows:

- 15 i. PDE I, rabbit heart
- ii. PDE II, rabbit heart
- iii. PDE III, rabbit heart, Jurkat cells
- iv. PDE IV, HL60 cells, rabbit brain, rabbit kidney and human recombinant PDE IV
- v. PDE V, rabbit lung, guinea pig lung

20

A gene encoding human PDE IV has been cloned from human monocytes (*Livi, et al., 1990, Molecular and Cellular Biology, 10, 2678*). Using similar procedures we have cloned human PDE IV genes from a number of sources including eosinophils, neutrophils, lymphocytes, monocytes, brain
25 and neuronal tissues. These genes have been transfected into yeast using an inducible vector and various recombinant proteins have been expressed which have the biochemical characteristics of PDE IV (*Beavo and Reifsnnyder, 1990, TIPS, 11, 150*). These recombinant enzymes, particularly the human eosinophil recombinant PDE IV, have been used as
30 the basis of a screen for potent, selective PDE IV inhibitors.

The enzymes were purified to isoenzyme homogeneity using standard chromatographic techniques.

35 Phosphodiesterase activity was assayed as follows. The reaction was conducted in 150µl of standard mixture containing (final concentrations):

50mM 2-[[tris(hydroxymethyl)methyl]amino]-1-ethane-sulphonic acid (TES)-NaOH buffer (pH 7.5), 10mM MgCl₂, 0.1μM [³H]-cAMP and vehicle or various concentrations of the test compounds. The reaction was initiated by addition of enzyme and conducted at 30°C for between 5 to 30 mins.

5 The reaction was terminated by addition of 50μl 2% trifluoroacetic acid containing [¹⁴C]-5'AMP for determining recovery of the product. An aliquot of the sample was then applied to a column of neutral alumina and the [³H]-cAMP eluted with 10ml 0.1 TES-NaOH buffer (pH8). The [³H]-5'-AMP product was eluted with 2ml 2M NaOH into a scintillation vial containing

10 10ml of scintillation cocktail. Recovery of [³H]-5'AMP was determined using the [¹⁴C]-5'AMP and all assays were conducted in the linear range of the reaction.

The salt according to the invention causes a concentration-dependent

15 inhibition of recombinant PDE IV at 0.1 - 1000nM with little or no activity against PDE I, II, III or V at concentrations up to 100μM.

2. The Elevation of cAMP in Leukocytes

The effect of the salt of the invention on intracellular cAMP was

20 investigated using human neutrophils or guinea pig eosinophils. Human neutrophils were separated from peripheral blood, incubated with dihydrocytochalasin B and the test compound for 10 min and then stimulated with FMLP. Guinea pig eosinophils were harvested by peritoneal lavage of animals previously treated with intra-

25 peritoneal injections of human serum. Eosinophils were separated from the peritoneal exudate and incubated with isoprenaline and test compound. With both cell types, suspensions were centrifuged at the end of the incubation, the cell pellets were resuspended in buffer and boiled for 10 min prior to measurement of cAMP by specific

30 radioimmunoassay (DuPont).

The salt according to the invention induced a concentration-dependent elevation of cAMP in neutrophils and/or eosinophils at a concentration of 0.1nM.

35

3. Suppression of Leukocyte Function

5 The salt of the invention were investigated for its effects on superoxide generation, chemotaxis and adhesion of neutrophils and eosinophils. Isolated leukocytes were incubated with dihydrocytochalasin B for superoxide generation only and test compound prior to stimulation with FMLP. The salt according to the invention caused a concentration-dependent inhibition of superoxide generation, chemotaxis and adhesion at a concentrationsof 0.1nM.

4. **Adverse Effects**

10 The salt according to the invention is free from adverse effects following repeated overdosage to rats or dogs.

CLAIMS

1. (R)-4-[2-(3-Cyclopentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine
hydrogen sulphate salt.
5
2. A pharmaceutical composition comprising (R)-4-[2-(3-Cyclo-
pentyloxy-4-methoxyphenyl)-2-phenylethyl]pyridine hydrogen
sulphate salt together with one or more pharmaceutically acceptable
carriers, excipients or diluents.
10
3. A process for the preparation of (R)-4-[2-(3-cyclopentyloxy-4-
methoxyphenyl)-2-phenylethyl]pyridine hydrogen sulphate salt which
comprises reacting (R)-4-[2-(3-cyclopentyloxy-4-methoxyphenyl)-2-
phenylethyl]pyridine and/or a salt thereof with sulphuric acid.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 95/01460

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D213/30 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO,A,94 14742 (CELLTECH LIMITED) 7 July 1994 cited in the application see claims 13,17; examples 3a,16i -----	1-3

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

20 September 1995

Date of mailing of the international search report

0 2. 10. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tlx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Bosma, P

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/GB 95/01460

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9414742	07-07-94	AU-B- 5709294	19-07-94
		CA-A- 2129039	07-07-94
		CN-A- 1092064	14-09-94
		CZ-A- 9402034	15-12-94
		EP-A- 0626939	07-12-94
		FI-A- 943856	22-08-94
		GB-A- 2279075	21-12-94
		JP-T- 7504442	18-05-95
		NO-A- 943091	21-10-94
		PL-A- 304892	09-01-95
